



Use for: Spotted fever rickettsiosis (SFR) including Rocky Mountain spotted fever (RMSF), Ehrlichiosis (*E. chaffeensis*, *E. ewingii*, & undet.), and Anaplasmosis (*A. phagocytophilum* & undet.). Visit <http://www.cdc.gov> and use "Search" for complete Case Definition(s) or visit the disease web site(s) for a fillable/downloadable PDF version of this Case Report.

1st COPY – STATE HEALTH DEPARTMENT



Visit <http://www.cdc.gov> and use "Search" for complete Case Definition(s) or visit the disease web site(s) for a fillable/downloadable PDF version of this Case Report.

Probable Ehrlichiosis/Anaplasmosis: A clinically compatible case with evidence of elevated IgG or IgM antibody reactive with *E. chaffeensis* or *A. phagocytophilum* antigen by IFA, enzyme-linked immunosorbent assay (ELISA), dot-ELISA, or assays in other formats (CDC uses an IFA IgG cutoff of $\geq 1:64$ and does not use IgM test results as independent diagnostic support criteria), OR identification of morulae in the cytoplasm of monocytes or macrophages (Ehrlichiosis) or in the cytoplasm of neutrophils or eosinophils (Anaplasmosis) by microscopic examination.

Public reporting burden of this collection of information is estimated to average 10 minutes per response. An agency may not conduct or sponsor, and a person is not required to respond to, a collection of information unless it displays a currently valid OMB control number. Please send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to CDC/ATSDR Reports Clearance Officer: 1600 Clifton Rd., NE (MS D-74): Atlanta, GA 30333; ATTN: PRA (0920-0009).

